

This is a repository copy of *Biodiversity and ecosystem services on the African continent – What is changing, and what are our options?*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/166814/>

Version: Accepted Version

---

**Article:**

Archer, E, Dziba, L E, Mulongoy, K J et al. (24 more authors) (2021) Biodiversity and ecosystem services on the African continent – What is changing, and what are our options? *Environmental Development*. 100558. ISSN 2211-4645

<https://doi.org/10.1016/j.envdev.2020.100558>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

## **Biodiversity and Ecosystem Services on the African continent – what is changing, and what are our options?**

Corresponding author: Prof Emma Archer, Centre for Environmental Studies/Geography, Geoinformatics and Meteorology, University of Pretoria cnr Lynnwood Road and Roper Street Hatfield 0028, South Africa. E-mail: [Emma.Archer@up.ac.za](mailto:Emma.Archer@up.ac.za)

Co authors:

L. E. Dziba, South African National Parks, 643 Leyds St, Muckleneuk, Pretoria, 0027, South Africa

K. J. Mulongoy, IEL, Montreal, Canada

M. A. Maoela, Council for Scientific and Industrial Research, Pretoria 0001, South Africa

M. Walters, Council for Scientific and Industrial Research, Pretoria 0001, South Africa

R. Biggs, Centre for Complex Systems in Transition (CST), University of Stellenbosch, Private Bag X1 Matieland 7602, South Africa

M-C. Cormier Salem, Institute of Research for Development, Marseille (IRD), France

F. DeClerck, EAT Foundation/Bioversity International, Parc Scientifique Agropolis II, 34397 Montpellier Cedex 5, France

M. C. Diaw, African Model Forest Network, Yaounde, Cameroon

A. E. Dunham, Department of Biosciences, Program in Ecology and Evolutionary Biology, Rice University, Houston, TX 77005, United States

P. Failler, University of Portsmouth, Richmond Building Portland Street Portsmouth P01 3DE, United Kingdom

C. Gordon, Institute for Environment and Sanitation Studies. College of Basic and Applied Sciences. University of Ghana, Legon, Accra, Ghana

K. A. Harhash, Biodiversity Central Department, Ministry of State for Environmental Affairs, Cairo, Egypt

R. Kasisi, Université de Montréal Faculté de l'aménagement, 2900 Edouard Montpetit Blvd, Montreal, Quebec H3T 1J4, Canada

F. Kizito, International Centre for Tropical Agriculture (CIAT), Kigali, Rwanda

W. D. Nyngi, Ichthyology Department, National Museums of Kenya, Nairobi, Kenya

N. Oguge, Centre For Advanced Studies in Environmental Law and Policy, University of Nairobi, Nairobi, Kenya

B. Osman-Elasha, African Development Bank, Abidjan, Côte d'Ivoire

L. C. Stringer, School of Earth and Environment, University of Leeds, Woodhouse Lane, Leeds, LS2 9JT, United Kingdom

L. Tito de Morais, Marine Environmental Sciences Laboratory, IRD, LEMAR Laboratory, Institut Universitaire Européen de la Mer, Technopôle Brest-Iroise, Rue Dumont D'urville 29280 Plouzané, France

A. Assogbadjo, Faculty of Agronomic Sciences, University of Abomey-Calavi, 05 BP 1752, Cotonou, Benin

B. N. Egoh, Earth System Science Department, University of California, Irvine, Croul Hall, Irvine, CA 92697-3100

M. W. Halmy, Department of Environmental Sciences Faculty of Sciences, Alexandria University Cairo, Egypt

K. Heubach, Palmengarten and Botanical Garden, Siesmayerstraße 61, 60323 Frankfurt am Main, Germany

A. Mensah, Institute for Environment and Sanitation Studies. College of Basic and Applied Sciences. University of Ghana, Legon, Accra, Ghana

L. Pereira, School of Arts and Social Sciences of City, University of London, Northampton Square, Clerkenwell, London EC1V 0HB, United Kingdom

N. Sitas, Centre for Complex Systems in Transition (CST), University of Stellenbosch, Private Bag X1 Matieland 7602, South Africa

Keywords: Biodiversity, ecosystem services, Africa, degradation, climate change

# **Biodiversity and Ecosystem Services on the African continent – what is changing, and what are our options?**

E. Archer, L. E. Dziba, K. J. Mulongoy, M. A. Maoela, M. Walters, R. Biggs, M-C. Cormier Salem, F. DeClerck, M. C. Diaw, A. E. Dunham, P. Failler, C. Gordon, K. A. Harhash, R. Kasisi, F. Kizito, W. D. Nyingi, N. Oguge, B. Osman-Elasha, L. C. Stringer, L. Tito de Morais, A. Assogbadjo, B. N. Egoh, M. W. Halmy, K. Heubach, A. Mensah, L. Pereira and N. Sitas

## **Abstract**

Throughout the world, biodiversity and nature's contributions to people are under threat, with clear changes evident. Biodiversity and ecosystem services have particular value in Africa– yet they are negatively impacted by a range of drivers, including land use and climate change. In this communication, we show evidence of changing biodiversity and ecosystem services in Africa, as well as the current most significant drivers of change. We then consider five plausible futures for the African continent, each underlain by differing assumptions. In three out of the five futures under consideration, negative impacts on biodiversity and ecosystem services are likely to persist. Those two plausible futures prioritizing environment and sustainability, however, are shown as the most likely paths to achieving long term development objectives without compromising the continent's biodiversity and ecosystem services. Such a finding shows clearly that achievement of such objectives cannot be separated from full recognition of the value of such services.

## **1. Introduction**

Biodiversity and ecosystem services are facing serious threats globally, impacted by a range of often interacting drivers, including land use and climate change (IPBES 2019). Africa, a continent rich in biocultural diversity, is one of the last places on Earth with a significant, intact large mammal assemblage, and with a unique diversity of indigenous and local knowledge, the majority of which, as yet, remains largely undocumented. The unrealized potential of Africa's biodiversity, ecosystem services, spirituality, culture and identities places the continent in a unique position globally- it can serve as a source for generating development pathways that are truly sustainable, where people's wellbeing and needs can be met without negatively infringing on the environment. The continent's rich biocultural heritage is, however, rapidly being exploited to meet development needs both within and outside of the continent. This has placed Africa in a vulnerable position with regards to building a resilient future for its citizens, and for those people and ecosystems that depend on Africa's resources outside the continent.

In this short communication, we draw on the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES) Regional Assessment Report on Biodiversity and Ecosystem Services for Africa – worked on by all authors. We show what is changing in biodiversity and ecosystem services on the African continent. We also identify future pathways and options for an African continent where long-term development objectives are recognized as inseparably connected to the conservation of the region's rich biocultural heritage.

## 2. . Material and approach

The Intergovernmental Science Policy Platform on Biodiversity and Ecosystem Services (IPBES) was established in 2012, with the intention of providing the most up to date and independent assessments of the state of biodiversity and ecosystem services (or nature's contributions to people) to support decision-making around the world. The Regional Assessment Report on Biodiversity and Ecosystem Services for Africa forms one of a suite of regional assessments, alongside those focusing on Asia-Pacific, Europe and Central Asia and the Americas, all of which were undertaken between 2015 and 2018.

The Africa Assessment was produced by 127 experts, including seven Fellows (early career scientists brought on at the start of the assessment); with support from 23 contributing authors. Authors were drawn largely from Africa. The report, as well as its Summary for Policymakers, was approved by the Member States of IPBES at the sixth session of the IPBES Plenary, in March 2018, in Medellín, Colombia.

## 3. What is changing?

Over the past several decades, biodiversity and ecosystem services in Africa have become increasingly threatened by anthropogenic drivers, some of the most important of which include human migration and political insecurity, climate change, habitat degradation and conversion, unsustainable harvesting and illegal trade of wildlife, and invasive alien species (MA, 2005; IPBES, 2018). Changes in land use and climate appear to be the most concerning of the drivers (more detail provided below); with land use change the primary driver of change and loss to date. Given current vulnerability to climate change in Africa (IPCC 2018), future changes in biodiversity and ecosystem services are likely to be exacerbated or driven by climate change, whether acting as a direct driver or in the case of multiple stressors. Natural drivers of biodiversity decline have also been increasing over the last two decades, including (but not limited to) diseases, pests and natural disasters (IPBES, 2018), likely as a result of human-driven environmental changes affecting the region (Daszak et al., 2000). Such increasing impacts have clear implications for a range of plants, invertebrates, fish, amphibians, reptiles, birds, mammals and micro-organisms (IPBES 2018).

Table 1 shows a qualitative assessment of change in intensity of drivers of change in biodiversity in Africa per sub-region and ecosystem type, as reported by parties to the Convention on Biological Diversity (CBD). We see here, for example, that climate change and habitat conversion are increasing in intensity, and may significantly impact both terrestrial/inland waters and coastal/marine biodiversity in all subregions.

Table 1: Changes in biodiversity and the role of underlying direct and indirect drivers in Africa shown per subregion and ecosystem type

**TABLE 1: CHANGES IN BIODIVERSITY AND THE ROLE OF UNDERLYING DIRECT AND INDIRECT DRIVERS IN AFRICA SHOWN PER SUBREGION AND ECOSYSTEM TYPE**

SUBREGIONS	ECOSYSTEM TYPE	DRIVERS OF BIODIVERSITY CHANGE							
		Direct drivers						Indirect drivers	
		Climate change	Habitat conservation	Overharvesting	Pollution	Invasive alien species	Illegal wildlife trade	Demographic change	Protected areas
CENTRAL AFRICA	Terrestrial/Inland waters	↗	↑	↑	↑	↑	↑	↑	↗
	Coastal/Marine	↗	↑	↑	↗	↗	↑	NI	↔
EAST AFRICA AND ADJACENT ISLANDS	Terrestrial/Inland waters	↑	↗	↑	↗	↗	↑	↑	↗
	Coastal/Marine	↑	↔	↗	↗	↗	↑	↑	↔
NORTH AFRICA	Terrestrial/Inland waters	↑	↗	↗	↗	↑	↔	→	→
	Coastal/Marine	↗	↗	↗	↗	↑	NI	→	→
SOUTHERN AFRICA	Terrestrial/Inland waters	↗	↗	↑	↗	↑	↗	↗	↗
	Coastal/Marine	↗	↗	↗	↗	↑	↗	↗	↗
WEST AFRICA	Terrestrial/Inland waters	↑	↑	↑	↗	↗	↑	↗	→
	Coastal/Marine	↑	↗	↗	↗	→	↑	↗	→

**DIRECTION OF ARROW:** Trend of the respective impact of the driver

High increase
 Moderate increase
 Low increase
 Decrease
 NI No information available
 Unchanged/Under control

It is well established that Africa is prone to the adverse impacts of climate change (see, for example, Myhre et al., 2013; Wright et al., 2015; Connolly-Boutin & Smit, 2016; Li et al., 2019). Temperatures throughout the continent are projected to rise more rapidly than the global rate

(IPBES, 2018; IPCC, 2018). In addition, there is a high probability that high intensity extreme rainfall events will increase in frequency (Akumaga & Tarhule, 2018). The most severe projections suggest that distribution, migration and population sizes of African plant species critical for food security (e.g., common bean) are likely to be affected by climate change (see Hummel et al., 2018). By 2100, it is estimated that climate change could result in significant loss of certain bird and mammal species (due to range retraction), and cause a decline in productivity of Africa's lakes by more than 20% (IPBES 2018).

In addition, climate change impacts on pests and pathogens are likely to significantly affect human health and the livestock sector throughout the continent (e.g., Bett et al., 2019; IPBES 2018). Negative climate change impacts on marine and coastal environments (e.g. salinization of water and soil, coastal erosion) pose a substantial risk for fisheries and the regulating and cultural ecosystem services these systems provide. For instance, extreme ocean warming caused massive coral bleaching events in 1998 and 2016, which resulted in reef mortality of more than 50% in certain regions (Obura, 2016), particularly the Western Indian Ocean (Gudka et al., 2018). Climate change and marine heatwaves (Smale et al., 2019), coupled with marine protected areas for which spatial data is available covering only 2.6% of Africa's marine jurisdiction (Belle et al., 2015), increases the impacts of current and future harvesting pressures on marine resources.

Land cover change throughout the continent is already driving a loss of key natural assets and reducing the continent's capacity to support biodiversity. Land cover change includes intensive agriculture, unregulated conversions of intact forest, mining, and use for urban and infrastructure development (IPBES 2018). Effectively, we are seeing the impact of competing demand for land through urban/infrastructure development, extractive industries and agricultural expansion and intensification – an example here would be development and investment choices that strongly emphasize expansion and intensification of primary and extractive industries. An estimate of 20 % of Africa's land surface is degraded due to direct drivers of change such as vegetation loss and adverse impacts on soils, including pollution, erosion, decreased fertility and salinization (Nyingi et al., 2018). In a significant finding, agricultural expansion appears as a dominant driver of biodiversity loss with unregulated conversion to agricultural land leading to loss and erosion of soils, habitats and water catchments, thus hampering Africa's long-term sustainable development (IPBES 2018). The interactions between land-use and climate change compound the impacts on biodiversity and ecosystem functioning, with ecosystems in environments that are climatically challenging displaying lowered resistance to land-use change (Peters et al., 2019).

Tackling the negative impact of these drivers of change is a critical aspect for sustainable development on the continent. Most African countries have committed to achieving particular targets by particular deadlines – including (but not limited to) the Aichi Biodiversity targets and the Sustainable Development Goals; as well as, for the continent specifically, AU Agenda 2063. Some countries are progressing well towards their targets and are on track within the mandated timeframe; others are not (Figure 1). For instance, awareness of biodiversity (Aichi Target 1) has grown, exceeding the target in some countries (Stringer et al 2018). For Aichi Target 10 which calls for reduction of pressures on ecosystems vulnerable to climate change however, evidence of progress is lacking. Information to monitor progress is absent for several countries, while in six nations, the direction of travel is away from the target.

FIGURE 1: COUNTRIES' PROGRESS TOWARDS SELECTED AICHI TARGETS




Figure 1 also shows some progress in the case of Aichi Target 11 – namely, protected areas (although this finding should be placed in the context that much progress in the case of targets is still only effected on paper – we discuss more in terms of conservation success stories below). Thirty-nine countries are progressing towards the target, albeit at an insufficient rate (Stringer et al., 2018).

Opportunities exist to learn from examples of better practice, including how we might be able to scale up approaches and initiatives worthy of replication. One key example here, shown in Box 1, is the West African Marine Protected Area Network that supports the growth and maintenance of Marine Protected Areas (MPAs) in West African countries (Failler et al. 2019).

#### Box 1: The West African Marine Protected Area Network

West African MPAs have been set-up initially for the protection of the fish biomass and/or certain emblematic species (turtles, manatees, birds, etc.). With the implementation of the National Determined Contribution in the context of the Paris Agreement on Climate Change, they further play the role of supplying key services for mitigation (blue carbon sequestration mainly) and for adaptation (coastal protection for instance). Overall, their habitats provide about 25% more regulating services than similar ones without special protection (Failler and Binet, 2012). A recent study, for example, showed that the Banc d'Arguin National Park, the largest African coastal MPA, would contribute to 20% of Mauritania's mitigation objective valued at 9 billion euros (with an annual running cost of only 1 million euros). Thus, the government, while recognising the key role of MPA, is taking steps toward the integration of coastal ecosystem services into its NDC (Tregarot et al., 2019). In other words, those measures put in place for the preservation of the biodiversity are now benefiting the society far beyond their initial mandate, with a very high return on public investment.

Indeed, as shown in Box 1 and elsewhere, protected areas serve as a key example of measures that are already contributing to the recovery of some threatened species. A further example here is the African Wild Dog (*Lycaon pictus*) in southern Africa (Davies-Mostert et al., 2009). Prudent land uses that maintain extensive, well-connected wildlife habitats, and reduce conflict with farmers through careful herding of livestock, have also been shown to facilitate recovery of the African wild dog in East Africa (Woodroffe, 2011), while Dube (2020) working in the Waterberg Biosphere Reserve in South Africa, highlights innovative measures for private landowners to monitor and track wild dogs, helping to reduce human-carnivore conflict. The example of the African Wild Dog is particularly interesting, since it includes land ownership and management that falls outside of, for example, formally designated national and provincial parks.

Other measures include control of alien invasive species and restoration of ecosystems (Nyingi et al., 2018), for example as articulated in the Volta Basin Authority's Strategic Action Plan. As at 2015, 13.4% of the continent's terrestrial and 2.6 % of the marine realm had been declared as protected areas (Belle et al., 2015); with other sites identified as wetlands of international importance, significant bird and biodiversity areas, community conserved areas, UNESCO World Heritage Sites, and Biosphere reserves, amongst others.








#### 4. Future pathways and options

Understanding the directions of changes to biodiversity and ecosystem services, and their contributions to human wellbeing can provide useful insights into how future changes could impact progress towards key targets, such as those outlined in the African Union Agenda 2063, the Sustainable Development Goals, and the post-2020 Aichi Biodiversity targets. The Africa Regional Assessment considered five plausible futures (Table 2) based on an archetype approach (Sitas and Harmáčková *et al.* in press) – all underpinned by various assumptions as to what each future could look like.

Table 2: The Global Scenarios Group (GSG) archetypes (at the global level) with their key characteristics and assumptions. Source: based on van Vuuren *et al.* (2012) (taken with permission from Biggs *et al.* 2018)

**TABLE 2: THE GLOBAL SCENARIOS GROUP (GSG) ARCHETYPES (AT THE GLOBAL LEVEL) WITH THEIR KEY CHARACTERISTICS AND ASSUMPTIONS**

GSG ARCHETYPE CATEGORY	 FORTRESS WORLD	 MARKET FORCES	 POLICY REFORM	 LOCAL SUSTAINABILITY	 REGIONAL SUSTAINABILITY
MAIN OBJECTIVES	Security	Economic growth	Various goals	Local sustainability	Regional and global sustainability
GLOBAL POPULATION GROWTH	High	Low	Low	Medium	Low
GLOBAL TECHNOLOGY DEVELOPMENT	Slow	Rapid	Rapid	Ranging from slow to rapid	Ranging from mid to rapid
GLOBAL ECONOMIC DEVELOPMENT	Slow	Very rapid	Rapid	Ranging from mid to rapid medium	Rangin from slow to rapid
TRADE	Trade barriers	Globalization	Globalization	Trade barriers	Globalization
POLICIES AND INSTITUTIONS	Strong national governments	Policies create open markets	Policies reduce market failures	Local steering: local actors	Strong global governance
ENVIRONMENTAL MANAGEMENT	Reactive	Reactive	Both reactive and proactive	Proactive	Proactive

The analysis showed that drivers of adverse changes in biodiversity and ecosystem services will increase under all the scenarios (Biggs *et al.* 2018). In turn, such changes are likely to further negatively impact on the ability of nature to contribute to human wellbeing and sustainable development under most cases, except in regional and local sustainability and supportive policy reform. It was unlikely that the African Union Agenda 2063, the SDGs and the Aichi Biodiversity would be achieved in three out of the five different futures (see Figure 2). Overall, only the regional and local sustainability futures offered pathways that offer Africa the greatest chances to meet its development goals in an economic, social and environmentally friendly way (Biggs *et al.* 2018).


All future scenarios present trade-offs but multiple synergies and policy alignments can support the feasibility of more desirable, equitable and sustainable development options. Our assessment

demonstrated that the 'Fortress World' scenario was least likely to support Africa in the achievement of multiple goals and targets. Overall, this future was found to result in failure to achieve important development goals. Market forces (MF) and policy reform (PR) scenarios, representing 'Business-as-usual' approaches, offer some potential for achieving multiple policy goals. Nevertheless, these futures do not adequately support biodiversity conservation, nor the diverse benefits of nature to human well-being. Conditions under a more 'managed transformation' type of future, through policies and practices aligned with regional sustainability and, to a lesser extent, local sustainability, increased the likelihood of reaching a range of sustainable goals.

Taking all the goals, targets and aspirations together, no single scenario option allows Africa to achieve them all, despite that some pathways appear more desirable for decision makers. If Africa is to achieve a desirable future (including that envisaged by commitment to targets), it is critical that development of policy and practice be not only based on inclusive and responsible economic tools, but also support the conservation and sustainable use of natural resources and their benefits to people (Figure 2).

208

209 Figure 2: Likelihood of achieving key outcomes in Africa under different global scenarios  
210 archetypes



211

212

213

214 **5. Conclusions: where to from here?**

215 As shown, there are options for Africa to balance development goals with protection of  
216 biodiversity and ecosystem services – in fact, such protection forms the basis for achieving  
217 development goals and improved human well-being. This may only be achieved, however,  
218 through a commitment to transformative change. Progress towards achievement of the Aichi  
219 Biodiversity Targets, SDGs, African Union's Agenda 2063, and the 2°C commitment under the  
220 2015 Paris Agreement on climate change, whilst helping support aspirations for a prosperous  
221 Africa, requires a fundamental shift away from the status quo.

Such transformative change towards sustainability, in line with aforementioned targets, will also depend on governance options that are able to harness synergies and deliver multiple benefits (IPBES 2018). By promoting policy coherence with adequate resources and capacity, and encouraging adaptive governance approaches that bring together different perspectives, a more equitable approach to accessing natural resources can ensue, helping to more effectively distribute costs and benefits. In addition, a more enabling environment that embraces Africa's diversity will help to ensure justice and fairness in access to the continent's diverse natural resources. A key finding here is that success stories regarding, for example, species stabilization or recovery, can not only rely on conservation within formal protected areas. This is, of course, a long addressed argument – but it is strongly emphasized in our review of those measures that might be scaled up. Measures that focus, for example, on private landowners or land managers outside of formally designated protected areas are clearly absolutely key (and must be evidence based). Africa has an ambitious development agenda that is critically tied to maintaining and sustainably harnessing its diverse natural systems, biodiversity and ecosystem services – as we have shown, they cannot be decoupled. In order to achieve this transformative agenda, it is necessary for all stakeholders to make use of effective policies that minimise trade-offs and maximise synergies under uncertainty so as to achieve a desirable and prosperous future for Africa.

We cannot conclude this paper without addressing COVID19, and the situation within which African conservation finds itself (this paper was first submitted in October 2019, and our context has, of course, changed dramatically). Certain models of conservation in Africa rely, to varying extents, on international tourism – and the recovery of this sector will be key to it's long term ability to achieve, for example, those biodiversity targets where regions and countries currently face difficulties (see, for example, Lindsey et al 2020 and their consideration of how to achieve conservation on the continent during COVID19, and in the post COVID19 period). In turn, conservation of biodiversity and ecosystem services is, of course, key to preventing and controlling zoonotic disease. As stated above, the continent has an ambitious development agenda – one that, along with the world at large, now faces possibly it's greatest economic challenge to date. To quote Lenzen et al in their recent paper in PLOSONe – 'How humanity reacts to this crisis will define the post pandemic world' (Lenzen et al 2020: 1). We can truly say that the post pandemic conservation world will help define our future, as a continent and as a planet.

## Acknowledgements

We would like to acknowledge the comments of two anonymous reviewers, as well as the editor – they were very useful. We would also like to thank Kerry Irish de Gale for work undertaken in redrawing our infographics, and in making them more understandable and accessible.

## References

- Akumaga, U., & Tarhule, A. (2018). Projected Changes in Intra-Season Rainfall Characteristics in the Niger River Basin, West Africa. *Atmosphere*, 9, 497. <https://doi.org/10.3390/atmos9120497>
- Belle, E., Wicander, S., Bingham, H., & Shi, Y. (2015). *Governance of Protected Areas in Africa, A Global Review*. UNEP-WCMC, Cambridge, UK. Available from <http://papaco.org/wp->

265 [content/uploads/2015/04/PAPACO-study-17\\_GOVERNANCE-STUDY-0-](content/uploads/2015/04/PAPACO-study-17_GOVERNANCE-STUDY-0-)  
266 [FINAL\\_REPORT-CONTEXT.pdf](#).

267 Bett, B., Lindahl, J., & Delia, G. (2019). Climate Change and Infectious Livestock Diseases: The  
268 Case of Rift Valley Fever and Tick-Borne Diseases. In The Climate-Smart Agriculture Papers:  
269 Investigating the Business of a Productive, Resilient and Low Emission Future. Edited by T.S.  
270 Rosenstock, A. Nowak, & E. Girvetz. Springer International Publishing, Cham. pp. 29–37.  
271 doi:10.1007/978-3-319-92798-5\_3.

272 Biggs, R., Kizito, F., Adjonou, K., Ahmed, M. T., Blanchard, R., Coetzer, K., Handa, C. O.,  
273 Dickens, C., Hamann, M., O’Farrell, P., Kellner, K., Reyers, B., Matose, F., Omar, K., Sonkoue,  
274 J-F., Terer, T., Vanhove, M., Sitas, N., Abrahams, B., Lazarova, T., & Pereira, L. Chapter 5:  
275 Current and future interactions between nature and society. In IPBES (2018): The IPBES  
276 regional assessment report on biodiversity and ecosystem services for Africa. Archer, E. Dziba,  
277 L., Mulongoy, K. J., Maoela, M. A., & Walters, M. (eds.). Secretariat of the Intergovernmental  
278 Science-Policy Platform on Biodiversity and Ecosystem Services, Bonn, Germany, pp. 297–352.

279 Connolly-Boutin, L., & Smit, B. (2016). Climate change, food security, and livelihoods in sub-  
280 Saharan Africa. Regional Environmental Change, 16(2), 385–399.  
281 <https://doi.org/10.1007/s10113-015-0761-x>

282 Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2000). Emerging infectious diseases of wildlife–  
283 threats to biodiversity and human health. Science, 287(5452), 443–449.  
284 <https://doi.org/10.1126/science.287.5452.443>

285 Davies-Mostert, H.T., Mills, M.G.L. & Macdonald, D.W. (2009). A critical assessment of South  
286 Africa’s managed metapopulation recovery strategy for African wild dogs. In Reintroduction of  
287 top-order predators. [Conservation Science and Practice No 5.]: 10–42. Hayward, M.W. &  
288 Somers, M.J. (Eds.).

289 Dube, L. (2020). Drivers of farmer-wild dog (lycaon pictus) conflict in the Waterberg Biosphere  
290 Reserve. MSc thesis, University of Pretoria. Pretoria, South Africa.

291 Failler P., Binet T. (2012), Évaluation de la valeur socio-économique des écosystèmes marins et  
292 côtiers des Aires marines protégées de l’Afrique de l’Ouest, Rapport synthèse,  
293 RAMP/PAO/PRCM/EVA, 8 p.

294 Failler, P., G. Touron-Gardic, M.-S. & Traoré (2019). Is Aichi Target 11 Progress correctly  
295 measured for developing Countries? Trends in Ecology and Evolution, available online 05 Aug.  
296 2019, <https://doi.org/10.1016/j.tree.2019.07.007>.

297 Hummel, M., Hallahan, B.F., Brychkova, G., Ramirez-Villegas, J., Guwela, V., Chataika, B.,  
298 Curley, E., McKeown, P.C., Morrison, L., Talsma, E.F., Beebe, S., Jarvis, A., Chirwa, R., &  
299 Spillane, C. (2018). Reduction in nutritional quality and growing area suitability of common bean  
300 under climate change induced drought stress in Africa. Sci Rep 8(1): 1–11. doi:10.1038/s41598-  
301 018-33952-4.

302 IPBES (2018). The IPBES regional assessment report on biodiversity and ecosystem services for  
303 Africa. Archer, E. Dziba, L., Mulongoy, K. J., Maoela, M. A., & Walters, M. (eds.). Secretariat of  
304 the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services, Bonn,  
305 Germany. 492 pages. <http://doi.org/10.5281/zenodo.3236178>

306 IPBES (2019). Summary for policymakers of the global assessment report on biodiversity and  
 307 ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and  
 308 Ecosystem Services. S. Díaz, J. Settele, E. S. Brondizio E.S., H. T. Ngo, M. Guèze, J. Agard, A.  
 309 Arneth, P. Balvanera, K. A. Brauman, S. H. M. Butchart, K. M. A. Chan, L. A. Garibaldi, K.  
 310 Ichii, J. Liu, S. M. Subramanian, G. F. Midgley, P. Miloslavich, Z. Molnár, D. Obura, A. Pfaff, S.  
 311 Polasky, A. Purvis, J. Razzaque, B. Reyers, R. Roy Chowdhury, Y. J. Shin, I. J. Visseren-  
 312 Hamakers, K. J. Willis, & C. N. Zayas (eds.). IPBES secretariat, Bonn, Germany.

313

314 IPCC (2018). Global warming of 1.5°C. An IPCC Special Report on the impacts of global  
 315 warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission  
 316 pathways, in the context of strengthening the global response to the threat of climate change,  
 317 sustainable development, and efforts to eradicate poverty [V. Masson-Delmotte, P. Zhai, H. O.  
 318 Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S.  
 319 Connors, J. B. R. Matthews, Y. Chen, X. Zhou, M. I. Gomis, E. Lonnoy, T. Maycock, M. Tignor,  
 320 T. Waterfield (eds.)].

321 Lenzen M., Li M., Malik A., Pomponi F., Sun Y-Y. & Wiedmann T., et al. (2020) Global socio-  
 322 economic losses and environmental gains from the Coronavirus pandemic. PLoS ONE 15(7):  
 323 e0235654. <https://doi.org/10.1371/journal.pone.0235654>

324 Li, L., Cao, R., Wei, K., Wang, W., & Chen, L. (2019). Adapting climate change challenge: A new  
 325 vulnerability assessment framework from the global perspective. Journal of Cleaner Production  
 326 217: 216–224. doi:10.1016/j.jclepro.2019.01.162.

327 Lindsey, P., Allan, J., Brehony, P., Dickman, A., Robson, A., Begg, C., Bhammar, H., Blanken,  
 328 L., Breuer, T., Fitzgerald, K. & Flyman, M. (2020). Conserving Africa's wildlife and wildlands  
 329 through the COVID-19 crisis and beyond. Nature Ecology & Evolution: 1-11.

330 MA (Millennium Ecosystem Assessment) (2005). Ecosystems and human well-being: Synthesis.  
 331 Washington, DC, USA: Island Press. Retrieved from <https://www.millenniumassessment.org/documents/document.356.aspx.pdf>

332

333 Gudka, M., Obura, D., Mwaura, J., Porter, S, Yahya, S. & Mabwa, R. (2018). Impact of the 3rd  
 334 Global Coral Bleaching Event on the Western Indian Ocean in 2016. Global Coral Reef  
 335 Monitoring Network (GCRMN)/Indian Ocean Commission. pp. 65

336 Myhre, G., Shindell, D., Bréon, F-M., Collins, W., Fuglestedt, J., Huang, J., Koch, D.,  
 337 Lamarque, J-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., &  
 338 Zhang, H. (2013). Anthropogenic and natural radiative forcing. In T. F. Stocker, D. Qin, G-K.  
 339 Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, & P. M. Midgley (Eds.),  
 340 Climate change 2013: The physical science basis. Contribution of Working Group I to the Fifth  
 341 Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge, UK:  
 342 Cambridge University Press. Retrieved from  
 343 [https://www.ipcc.ch/pdf/assessmentreport/ar5/wg1/WG1AR5\\_Frontmatter\\_FINAL.pdf](https://www.ipcc.ch/pdf/assessmentreport/ar5/wg1/WG1AR5_Frontmatter_FINAL.pdf)

344 Nyingi, W., Ouge, N., Dziba, L., Chandipo, R., Didier, T. A., Gandiwa, E., Kasiki, S., Kisanga,  
 345 D., Kgosikoma, O., Osano, O., Tassin, J., Sanogo, S., von Maltitz, G., Ghazi, H., Archibald, S.,  
 346 Gambiza, J., Ivey, P., Logo, P. B., Maoela, M. A., Ndarana, T., Ogada, M., Olago, D., Rahlao, S.,  
 347 & van Wilgen, B. Chapter 4: Direct and indirect drivers of change in biodiversity and nature's  
 348 contributions to people. In IPBES (2018): The IPBES regional assessment report on biodiversity



349 and ecosystem services for Africa. Archer, E., Dziba, L., Mulongoy, K. J., Maoela, M. A., &  
 350 Walters, M. (eds.). Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity  
 351 and Ecosystem Services, Bonn, Germany, pp. 207–296.

352 Obura, D. O. (2016). Coral bleaching response guide 2016 (Western Indian Ocean). Retrieved  
 353 from [http://cordioea.net/bleaching\\_resilience/wio-bleaching-2016](http://cordioea.net/bleaching_resilience/wio-bleaching-2016)

354 Peters, M. K., Hemp, A., Appelhans, T., Becker, J. N., Behler, C., Classen, A., ... & Gebert, F.  
 355 (2019). Climate–land-use interactions shape tropical mountain biodiversity and ecosystem  
 356 functions. Nature, 568(7750), 88.

357 Smale, D. A., Wernberg, T., Oliver, E. C., Thomsen, M., Harvey, B. P., Straub, S. C., ... & Feng,  
 358 M. (2019). Marine heatwaves threaten global biodiversity and the provision of ecosystem  
 359 services. Nature Climate Change, 9(4), 306.

360 Stringer, L. C., Osman-Elasha, B., DeClerck, F., Ayuke, F. O., Gebremikael, M. B., Barau, A. S.,  
 361 Denboba, M. A., Diallo, M., Molua, E. L., Ngenda, G., Pereira, L., Rahlao, S. J., Kalemba, M. M.,  
 362 Ojino, J. A., Belhabib, D., Sitas, N., Strauß, L., & Ward, C. Chapter 6: Options for governance  
 363 and decision-making across scales and sectors. In IPBES (2018): The IPBES regional assessment  
 364 report on biodiversity and ecosystem services for Africa. Archer, E. Dziba, L., Mulongoy, K. J.,  
 365 Maoela, M. A., & Walters, M. (eds.). Secretariat of the Intergovernmental Science-Policy  
 366 Platform on Biodiversity and Ecosystem Services, Bonn, Germany, pp. 353–414.

367 Tambling, C. J., & Toit, J. T. D. (2005). Modelling Wildebeest Population Dynamics:  
 368 implications of predation and harvesting in a closed system. Journal of Applied Ecology, 42  
 369 (2005), pp. 431–441.

370 Trégarot E., Catry T., Pottier A., Cornet C., Maréchal J.-P., Fayad V., Sidi Cheick M.A., David G.,  
 371 Dia A.D., Fall A.D., Sarr O., El Valy Y., Wagne O.H., Meisse B., Kane E.A., Ball A.C., Haidallah  
 372 M.S., Braham C.B., Dia M., Abdel Hamid M.L., Rey-Valette H., Salles J.-M., Ly D., Gascuel D.,  
 373 Cissé C.B. & Failler P. (2018). Évaluation des services écosystémiques du Banc d'Arguin,  
 374 Mauritanie. Rapport final pour le Parc National du Banc d'Arguin, 402 p.

375 UNEP (2016). Blue Carbon Financing of Mangrove Conservation in the Abidjan Convention  
 376 Region: A Feasibility Study. United Nations Environment Programme, Abidjan Convention  
 377 Secretariat and GRID-Arendal, Nairobi, Abidjan and Arendal.

378 Van Vuuren, D.P., Kok, M.T., Girod, B., Lucas, P.L. & de Vries, B. (2012). Scenarios in global  
 379 environmental assessments: key characteristics and lessons for future use. Global Environmental  
 380 Change, 22(4), pp.884–895.

381 Woodroffe, R. (2011). Demography of a recovering African wild dog (*Lycaon pictus*)  
 382 population. Journal of Mammalogy, 92(2): 305–315

383 Wright, H., Huq, S., & Reeves, J. (2015). Impact of climate change on least developed countries:  
 384 Are the SDGs possible? Retrieved from <http://pubs.iied.org/17298IIED/>